



Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

Model Answers

Forename(s)

Candidate signature

AS CHEMISTRY

Paper 2 Organic and Physical Chemistry

Friday 25 May 2018

Morning

Time allowed: 1 hour 30 minutes

Materials

For this paper you must have:

- the Periodic Table/Data Sheet, provided as an insert (enclosed)
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do **not** write outside the box around each page or on blank pages.
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

Advice

- You are advised to spend about 65 minutes on **Section A** and 25 minutes on **Section B**.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
Section B	
TOTAL	

Section A

Answer **all** questions in this section.

0 1

Hydrogen peroxide solution decomposes slowly to form water and oxygen.
The reaction is much faster in the presence of a manganese(IV) oxide catalyst.




Three experiments, shown in **Table 1**, were carried out to investigate how the volume of oxygen produced varied over time under different conditions. The same mass of catalyst was used in each experiment.

Table 1

Experiment	Concentration of $\text{H}_2\text{O}_2(\text{aq})$ / mol dm^{-3}	Volume of $\text{H}_2\text{O}_2(\text{aq})$ / cm^3	Temperature / $^{\circ}\text{C}$	Catalyst
1	1.0	50	20	lumps
2	1.0	50	20	powder
3	0.5	50	20	lumps

Figure 1 shows how the volume of oxygen collected varied with time in Experiment 1.

Figure 1



0 1 . 1

Explain, in general terms, how a catalyst increases the rate of a reaction.

[2 marks]

A catalyst speeds up reactions by providing an alternate pathway which has a lower activation energy.

0 1 . 2

Draw **two** lines on **Figure 1** to show how the volume of oxygen collected varied with time in Experiments **2** and **3**.

Label each line with the experiment number.

[2 marks]

see graph.

0 1 . 3

Explain, in terms of collision theory, the effect of increasing the concentration of hydrogen peroxide on the rate of reaction.

[2 marks]

with increased concentration there are more particles of H_2O_2 in a given volume. This means there are more frequent collisions.

Interesting that Mark scheme
suggests more successful but for
conc. frequency is better.




0 2

Citric acid, $\text{C}_3\text{H}_5\text{O}(\text{COOH})_3$, occurs naturally in many fruits and can also be synthesised in the laboratory for use as a food flavouring. A student analysed a sample of citric acid to determine its percentage purity.

The student dissolved 784 mg of impure citric acid in water to prepare 250 cm^3 of solution in a volumetric flask.

The student titrated 25.0 cm^3 samples of this solution with $0.0500 \text{ mol dm}^{-3}$ sodium hydroxide solution using phenolphthalein as the indicator.



0 2 . 1

The student rinsed the burette before filling it with the sodium hydroxide solution.

State why the student should use sodium hydroxide solution rather than water for the final rinse of the burette.

[1 mark]

By using water you would effectively be diluting the sodium hydroxide.

0 2 . 2

The student carried out several titrations. The results are shown in **Table 2**.

Complete **Table 2** to show the titre in each titration.

[1 mark]

Table 2

Titration	Rough	1	2	3
Final reading / cm^3	25.2	23.95	47.65	24.10
Start reading / cm^3	0.0	0.05	23.95	0.10
Titre / cm^3	25.2	23.90	23.70	24.00

0 2 . 3

Calculate the mean titre using the concordant results.
Give your answer to the appropriate number of significant figures.

[2 marks]

$$\frac{23.90 + 24.00}{2} = 23.95 \text{ cm}^3$$

Mean titre _____ cm^3



0 2 . 4

The total uncertainty when using the burette is $\pm 0.15 \text{ cm}^3$. This is the combination of uncertainties in the start reading, final reading and the determination of the end point.

Use your answer to Question 02.3 to calculate the percentage uncertainty for the use of the burette in this experiment.

$$\left(\frac{0.15}{23.95} \right) \times 100 = 0.63\% \quad \left(\frac{\text{uncertainty}}{\text{titre}} \right) \times 100 \quad [1 \text{ mark}]$$

Percentage uncertainty _____ %

0 2 . 5

Use your answer to Question 02.3 to find the mass, in mg, of citric acid dissolved in 250 cm^3 of the solution.

The relative molecular mass (M_r) of citric acid is 192.0

$$\text{Conc} = \frac{\text{mol}}{\text{Vol}} \quad \text{Conc} \times \text{Vol} = \text{mol} \quad 0.05 \times \frac{23.95}{1000} = 1.1975 \times 10^{-3} \quad [3 \text{ marks}]$$

moles of NaOH in 25 cm^3 .

mole ratio from balanced equation. \rightarrow

$$\frac{1.1975 \times 10^{-3}}{3} = 3.99 \times 10^{-4} \text{ moles of citric acid in } 25 \text{ cm}^3 \quad \times 10 = 3.99 \times 10^{-3} \text{ citric acid moles in } 250 \text{ cm}^3$$

$$\text{moles} \times M_r = \text{mass}$$

$$3.99 \times 10^{-3} \times 192.0 = 0.76608 \quad \times 1000 = 766.08 \text{ mg}$$

$\text{g} \rightarrow \text{mg}$

or 766 mg

Mass _____ mg

0 2 . 6

Calculate the percentage purity of this sample of citric acid.

$$\frac{766}{784} \times 100 = 97.7\%$$

Percentage purity _____ %




0 3

This question is about enthalpy changes.

0 3

1

When ethanoic acid reacts with sodium hydroxide, the enthalpy change, ΔH , is $-56.1 \text{ kJ mol}^{-1}$



Calculate the temperature rise when 25 cm^3 of 2.0 mol dm^{-3} aqueous ethanoic acid react with 25 cm^3 of 2.0 mol dm^{-3} aqueous sodium hydroxide.

Assume that both solutions have the same initial temperature, have a density of 1.0 g cm^{-3} and a specific heat capacity of $4.18 \text{ J K}^{-1} \text{ g}^{-1}$

[4 marks]

$$Q = m \times c \times \Delta T$$

$$C \times V = \text{moles} \quad 2.0 \times \left(\frac{25}{1000}\right) = 0.05 \text{ moles of } \text{CH}_3\text{COOH} \quad \text{and } 0.05 \text{ moles of } \text{NaOH}$$

$$\Delta H = \frac{Q}{\text{mole}}$$

$$\Delta H \times \text{moles} = Q \quad 56.1 \times 0.05 = 2.805 \text{ kJ}$$

$$2.805 \times 1000 = 2805 \text{ J}$$

$$\Delta T = \frac{Q}{m \times c}$$

no need for negative as energy NOT enthalpy.

$$\Delta T = \frac{2805}{50 \times 4.18}$$

$$\Delta T = \underline{\underline{13.4^\circ\text{C}}}$$

total volume
of $50 \text{ cm}^3 = 50 \text{ g}$

Temperature rise _____ $^\circ\text{C}$



03.2

A student recorded the temperature of aqueous ethanoic acid in a polystyrene cup for three minutes.

At the fourth minute, the student added sodium hydrogencarbonate.

The student stirred the mixture and carried on recording the temperature every minute for several minutes.


The student's measurements are shown in **Figure 2**.

A best-fit line showing the temperature before mixing has been drawn.

Draw an appropriate best-fit line on **Figure 2** and use it to find the temperature change at the time of mixing.

[2 marks]

Figure 2




Temperature change at time of mixing 5.7 °C

0 4

The alkanes nonane and 2,4-dimethylheptane are structural isomers with the molecular formula C_9H_{20} .


They are found in crude oil and can be separated by fractional distillation. Both can be used in fuels or cracked to form other products.

nonane



boiling point 151 °C

2,4-dimethylheptane



boiling point 134 °C

0 4 . 1

State the general formula of an alkane containing n carbon atoms.

Deduce an expression for the relative molecular mass (M_r) of an alkane in terms of n .

[2 marks]

General formula



if $C=12$ and $H=1 \therefore 12 + 2H's = 14$

Expression

$$14n + 2$$

0 4 . 2

Explain why nonane has a higher boiling point than 2,4-dimethylheptane.

[2 marks]

Nonane would have more van der Waals forces between molecules as there would be more ordered packing due to the straight chain alkane having no branching.



0 4 . 3 Give an equation for the complete combustion of nonane.

[1 mark]



0 4 . 4 Nonane is often found in fuel for jet engines. Combustion in jet engines produces pollutants including nitrogen monoxide (NO).

Explain how this nitrogen monoxide is formed.

[2 marks]

Nitrogen and oxygen in the air are able to react due to the high temperatures in the engine.

0 4 . 5 Nonane can be cracked to form large quantities of propene.

Name the type of cracking used.


thermal gives high % of alkenes.
catalytic gives more aromatic + branched products [1 mark]

thermal cracking.

0 4 . 6 The main use of propene, formed from cracking, is to make poly(propene).

Draw the repeating unit of poly(propene).

[1 mark]



Turn over for the next question

- 0 5 . 1 A hydrocarbon contains 87.8% by mass of carbon and has a relative molecular mass (M_r) of 82.0
The hydrocarbon decolourises bromine water.

- (1) Determine the empirical and molecular formulae of the hydrocarbon.
(2) Suggest **two** possible structures for the hydrocarbon.
(3) Name the type of reaction taking place when bromine water reacts with the
(4) hydrocarbon.

[6 marks]

$$\frac{87.8}{12} = 7.316 \quad \frac{12.2}{1} = 12.2$$

$$\frac{7.316}{7.316} = 1 \quad \frac{12.2}{7.316} = 1.667$$


 $\times 3$

= 3 : 5


lowest whole
number ratio C_3H_5 = empirical formula (1)

If $C_3H_5 = 41 \therefore \times 2 = 82 \quad C_6H_{10}$ = molecular formula (2)

(3)




or



(4)


Electrophilic addition



0 6

Compound A is a halogenoalkane.

Compound A



0 6 . 1

Name Compound A.

→ alphabetical so chloro before methyl.

[1 mark]

3-chloro-2,4-dimethyl-pentane


0 6 . 2

Compound A has a relative molecular mass (M_r) of 134.5The main isotope of hydrogen is ^1H The main isotope of carbon is ^{12}C Chlorine consists of two common isotopes, ^{35}Cl and ^{37}Cl , of which 75% is ^{35}Cl The mass spectrum of A was recorded when A was ionised by electron impact to form A^+ ions.

Draw, on Figure 3, the peaks for the main molecular ions in the mass spectrum of A.

[2 marks]

Figure 3



*IF Cl atom is 35 then 134 but 3/1 ratio.
if Cl atom is 37 then 136*

Question 6 continues on the next page


0 6 . 3 Reaction of A with warm, dilute aqueous sodium hydroxide forms alcohol B.

- ① Name the mechanism for this reaction.
- ② Outline the mechanism using the structure of A shown.
- ③ Include the structure of the product, alcohol B.

[4 marks]

Mechanism Nucleophilic substitution

Outline of mechanism



0 6 . 4 Reaction of **A** with hot, ethanolic potassium hydroxide gives alkene **C**.


- 1 Name the mechanism for this reaction.
- 2 State the role of the hydroxide ions.
- 3 Outline the mechanism using the structure of **A** shown.
Include the structure of the product, alkene **C**.

[6 marks]

Mechanism Elimination

Role of hydroxide ions Base


Outline of mechanism



Question 6 continues on the next page

Turn over ►

06.5

The infrared spectrum in **Figure 4** is that of either alcohol **B** or alkene **C**.**Figure 4**

Tick the box that shows the correct compound.

Explain your answer with reference to a bond and the wavenumber of its absorption.

[1 mark]



Alcohol **B**☐Alkene **C**☒

Explanation

It is not alcohol B as there is no O-H peak(a) 3230-3550 cm⁻¹.Could have said C=C peak(a) 1620-1680 is presentfor alkene.

0 6 . 6

Compound **D** reacts with dilute aqueous sodium hydroxide in a similar way to **A** to form alcohol **B**.

Compound **A**Compound **D**

Explain why **D** reacts more quickly than **A** with dilute aqueous sodium hydroxide at the same temperature.

[1 mark]

C-Br is weaker than C-Cl so breaks more easily.

15

Turn over for the next question

Turn over ►



0 7 . 1 Four compounds, all colourless liquids, are

- butan-2-ol
- butanal
- butanone
- 2-methylpropan-2-ol

Two of these compounds can be identified using different test-tube reactions.

Describe these **two** test-tube reactions by giving reagents and observations in each case.

Suggest how the results of a spectroscopic technique could be used to distinguish between the **other** two compounds.


[6 marks]

- ① firstly use tollens reagent which would give a silver mirror with butanal but would have no reaction with the other three.
- ② Now use acidified potassium dichromate which would change colour from orange to green for both butanal and butan-2-ol but would not react with the other two.
- ③ We could distinguish between butanone and 2-methylpropan-2-ol by using I.R spectroscopy. Butanone would give a carbonyl ($C=O$) peak @ $1680-1750\text{cm}^{-1}$, but 2-methylpropan-2-ol would not. Instead it would give a peak @ $3230-3550\text{cm}^{-1}$ for the O-H alcohol group.



0 8

Methanol can be manufactured in a reversible reaction as shown by the equation.



0 8 . 1

State and explain the effect of using a catalyst on the yield of methanol in this equilibrium.

[2 marks]

A catalyst has no effect on the yield as it increases the rate of reaction in both the forward and backwards directions.

0 8 . 2

Give an expression for the equilibrium constant (K_c) for this reaction.

[1 mark]

$$K_c = \frac{[\text{CH}_3\text{OH}]}{[\text{CO}] [\text{H}_2]^2}$$



08.3

A mixture of carbon monoxide and hydrogen was allowed to reach equilibrium in a container of volume 250 cm^3 at temperature T .

At equilibrium, the mixture contained 0.340 mol of carbon monoxide, 0.190 mol of hydrogen and 0.0610 mol of methanol.

Calculate the value of the equilibrium constant (K_c) for this reaction at temperature T .

[3 marks]

$$K_c = \frac{[\text{CH}_3\text{OH}]}{[\text{CO}][\text{H}_2]^2}$$

$$K_c = \frac{0.0610}{\frac{(250/1000)}{0.340} \times \left(\frac{0.190}{(250/1000)}\right)^2}$$

Don't forget the
to turn them into
concentrations
so $\div \text{Vol!}$

$$K_c = \frac{0.244}{1.36 \times 0.5776}$$

$$K_c = \underline{\underline{0.311}}$$

$$K_c \text{ } \underline{\hspace{2cm}} \text{ mol}^{-2} \text{ dm}^6$$

08.4

Methanol decomposes on heating in a reaction that is the reverse of that used in its manufacture.



reverse equation
 \therefore reciprocal of the
above K_c .

Use your answer from Question 08.3 to determine the value of K_c for this equilibrium at temperature T .

State the units for this value of K_c .

(If you were unable to complete the calculation in Question 08.3, assume a value of $K_c = 0.825 \text{ mol}^{-2} \text{ dm}^6$. This is **not** the correct value.)

[2 marks]

$$\frac{1}{0.311} = \underline{\underline{3.22}}$$

Value of K_c 3.22

Units of K_c $\text{mol}^2 \text{ dm}^{-6}$

$\frac{1}{\text{mol}^{-2} \text{ dm}^6} \rightarrow$

Section B

Answer **all** questions in this section.Only **one** answer per question is allowed.

For each answer completely fill in the circle alongside the appropriate answer.

CORRECT METHOD



WRONG METHODS



If you want to change your answer you must cross out your original answer as shown.



If you wish to return to an answer previously crossed out, ring the answer you now wish to select as shown.

You may do your working in the blank space around each question but this will not be marked.
Do **not** use additional sheets for this working.

0 9

A student has a 10 cm^3 sample of $1.00 \times 10^{-2} \text{ mol dm}^{-3}$ methanoic acid solution. The student is asked to dilute the methanoic acid solution to a concentration of $2.00 \times 10^{-4} \text{ mol dm}^{-3}$ by adding distilled water.

Which volume of water should be added?

[1 mark]

A 200 cm^3 B 490 cm^3 C 500 cm^3 D 510 cm^3

$$C = \frac{n}{V} \quad 1 \times 10^{-2} = \left(\frac{10}{1000} \right) : 0.0001 \text{ moles}$$

$$\frac{n}{C} = V \quad \frac{0.0001}{2 \times 10^{-4}} = 0.5 \text{ dm}^3 \text{ or } 500 \text{ cm}^3$$


$\therefore 490$ needed to be added!



1 0

Which molecule does **not** have a permanent dipole?






[1 mark]

A CH_3Br B CH_2Br_2 C CHBr_3 D CBr_4 

1 1

Which is the major product of the reaction between 2-methylbut-2-ene and iodine monochloride (ICl)?

[1 mark]

☐☐☒☐

1 2

Which statement is **not** correct about the industrial preparation of ethanol by the hydration of ethene at 300 °C?



[1 mark]

A The reaction is catalysed by an acid.

☐

B The higher the pressure, the higher the equilibrium yield of ethanol.

☐

C The higher the temperature, the higher the equilibrium yield of ethanol.

☒

x at high temps eq will shift left to counteract the change.

D A low equilibrium yield of ethanol is acceptable because unreacted ethene is recycled.

☐

1 3 Which compound has the highest boiling point?

[1 mark]

- A butanal *Dipole-dipole*
- B butan-2-ol *Hydrogen bonding.*
- C but-2-ene *van der Waals*
- D 1-fluorobutane *dipole-dipole.*

☐
☒
☐
☐

1 4 Which statement is correct about the fractional distillation of crude oil?

[1 mark]

- A A zeolite catalyst is used.
- B Each fraction contains a mixture of hydrocarbons.
- C Gaseous fractions are formed by breaking covalent bonds.
- D The fractionating column is hottest at the top.

☐
☒
☐
☐

- no fraction is perfectly pure - mixture of similar fractions.

1 5 How many structural isomers with an unbranched carbon chain have the molecular formula $C_4H_8Br_2$?


[1 mark]

A 4

B 5

C 6

D 7



☐
☐
☒
☐

1 6

The Maxwell-Boltzmann distribution of molecular energies in a sample of gas at a fixed temperature is shown.

Which letter represents the mean energy of the molecules?

[1 mark]



A ☒ } not correct axis.

B ☒

C - most probable energy.

D - mean energy (always to right of maximum)

☐
☐
☐
☒

1 7

Ethanol can be made from glucose by fermentation.



In an experiment, 268 g of ethanol ($M_r = 46.0$) were made from 1.44 kg of glucose ($M_r = 180.0$).

What is the percentage yield?

[1 mark]

A 18.6%

B 36.4%

C 51.1%

D 72.8%

1440g of glucose = 1.44kg.

$$\text{moles} = \frac{\text{mass}}{M_r} \quad \frac{1440}{180} = 8 \text{ moles}$$

$$8 \times 2 = \text{moles of ethanol} \quad 16 \times 46 = 736 \text{g of ethanol}$$

$$\frac{268}{736} \times 100 = \underline{\underline{36.4\%}}$$

☐
☒
☐
☐

1 8 Which species could act as a nucleophile?

[1 mark]

A BH_3

☐

B NH_4^+

☐

C PH_3 *$\text{H}-\ddot{\text{P}}-\text{H}$ Lone pair available.*

☒

D SiH_4

☐

1 9 Which statement is correct about poly(chloroethene)?

[1 mark]

A It has the empirical formula CHCl *Not true*

☐

B It decolourises bromine water. *Not an alkene*

☐

C Its brittleness is reduced by plasticisers. *✓*

☒

D Its polymer chain contains alternate single and double bonds. *not true.*

☐

2 0 What is the enthalpy of formation of buta-1,3-diene, $\text{C}_4\text{H}_6(\text{g})$?

Substance	Enthalpy of combustion / kJ mol^{-1}
$\text{C}_4\text{H}_6(\text{g})$	-2546
$\text{C}(\text{s})$	-394
$\text{H}_2(\text{g})$	-286

[1 mark]

A $+112 \text{ kJ mol}^{-1}$

☒


B -112 kJ mol^{-1}

☐

C $+746 \text{ kJ mol}^{-1}$

☐

D -746 kJ mol^{-1}

☐


- 2 1 A gas cylinder contains 5.0 kg of propane. $\rightarrow C_3H_8 = 44$

How many propane molecules are in the cylinder?

The Avogadro constant, $L = 6.022 \times 10^{23} \text{ mol}^{-1}$

[1 mark]

A 6.8×10^{22}

B 7.2×10^{22}

C 6.8×10^{25}

D 7.2×10^{25}

$5 \text{ kg} = 5000 \text{ g}$

$\frac{5000}{44} = 113.64 \text{ moles}$

$\times 6.02 \times 10^{23}$

$= 6.84 \times 10^{25}$

☐
☐
☒
☐

- 2 2 Which sample of liquid has the greatest volume?

[1 mark]

A 500 mg of pentane (density = 0.63 g cm^{-3})

$\frac{0.5}{0.63} = 0.79$

B 650 mg of propan-1-ol (density = 0.80 g cm^{-3})

$\frac{0.65}{0.8} = 0.813$

C 1.20 g of dichloromethane (density = 1.33 g cm^{-3})

$\frac{1.2}{1.33} = 0.9$

D 1.30 g of trichloromethane (density = 1.48 g cm^{-3})



$\frac{1.3}{1.48} = 0.88$

$d = \frac{m}{v}$ $\frac{m}{d} = v$

☐
☐
☒
☐

- 2 3 Which equation represents an initiation step?

[1 mark]


☐
☐
☐
☒

END OF QUESTIONS

