

# F321 Atoms, Bonds and Groups



Question		Expected Answers	Marks	Additional Guidance
1	(a)	<p>Mass of the <b>isotope</b> compared to 1/12th  <b>OR</b>  <b>mass</b> of the <b>atom</b> compared to 1/12th ✓</p> <p>(the mass of a) carbon-12 <b>OR</b> <math>^{12}\text{C}</math> (atom) ✓</p>	2	<p><b>IGNORE</b> Reference to average <b>OR</b> weighted mean            (i.e. correct definition of relative atomic mass will score both marks)</p> <p><b>ALLOW</b> mass of a <b>mole</b> of the isotope/atom with 1/12th the mass of a <b>mole OR 12 g</b> of carbon-12 for two marks.</p> <p><b>ALLOW 2 marks for:</b>            'Mass of the isotope <b>OR</b> mass of the atom compared to <math>^{12}\text{C}</math> atom given a mass of 12.0'            i.e. 'given a mass of 12' <b>OR</b> C12 is 12 communicates the same idea as 1/12th.'</p> <p><b>ALLOW</b> 12C <b>OR</b> C12</p> <p><b>ALLOW 2 marks for:</b>  <math display="block">\frac{\text{mass of the isotope}}{\text{mass of 1/12th mass of carbon - 12}}</math>           i.e. fraction is equivalent to 'compared to'</p> <p><b>ALLOW 1 mark for</b> a mix of mass of atom and mass of mole of atoms, i.e. 'mass of the isotope/mass of an atom compared with 1/12th the mass of a <b>mole OR 12 g</b> of carbon-12.'</p> <p><b>DO NOT ALLOW</b> mass of 'ions' <b>OR</b> mass of element</p>
	(b)	<p><math display="block">\frac{(151 \times 47.77) + (153 \times 52.23)}{100}</math></p> <p><b>OR</b>            72.1327 + 79.9119  <b>OR</b>            152.0446 (calculator value) ✓  <math>A_r = 152.04</math> ✓</p>	2	<p><b>ALLOW</b> Correct answer for two marks</p> <p><b>ALLOW</b> One mark for ECF from transcription error in first sum provided final answer is to 2 decimal points and is to between 151 and 153 and is a correct calculation of the transcription</p>


Question		Expected Answers	Marks	Additional Guidance
	(c) (i)	$^{153}\text{Eu}$ has (2) more neutrons <b>OR</b> $^{153}\text{Eu}$ has 90 neutrons <b>AND</b> $^{151}\text{Eu}$ has 88 neutrons ✓	1	<b>ALLOW</b> There are a different number of neutrons <b>IGNORE</b> Correct references to protons / electrons <b>DO NOT ALLOW</b> Incorrect references to protons / electrons
	(ii)	(It has the) same number of protons <b>AND</b> electrons <b>OR</b> Both have 63 protons and 63 electrons ✓	1	<b>ALLOW</b> Same number of protons <b>AND</b> same electron configuration <b>DO NOT ALLOW</b> 'Same number of protons' without reference to electrons (and vice versa)

Question	Expected Answers	Marks	Additional Guidance
(d)	<p>Xe has a bigger atomic radius <b>OR</b> Xe has more shells ✓</p> <p>Xe has <b>more</b> shielding ✓</p> <p>The nuclear attraction decreases  <b>OR</b> Outermost electrons of Xe experience less attraction (to nucleus)  <b>OR</b> Increased shielding / distance outweighs the increased nuclear charge ✓  ORA throughout</p>	3	<p><b>ALLOW</b> Xe has more energy levels  <b>ALLOW</b> Xe has electrons in higher energy level  <b>ALLOW</b> Xe has electrons further from nucleus  <b>IGNORE</b> Xe has more orbitals <b>OR</b> more sub-shells  <b>DO NOT ALLOW</b> 'different shell' or 'new shell'</p> <p><b>ALLOW More</b> screening  There must be a clear comparison ie <b>more</b> shielding <b>OR</b> <b>increased</b> shielding.  i.e. <b>DO NOT ALLOW</b> Xe 'has shielding'  <b>ALLOW</b> Xe has <b>more</b> electron repulsion from inner shells</p> <p><b>ALLOW</b> Xe has less nuclear pull  <b>IGNORE</b> Xe has less effective nuclear charge  <b>DO NOT ALLOW</b> nuclear charge for nuclear attraction</p>
	<b>Total</b>	<b>9</b>	

Question			Expected Answers	Marks	Additional Guidance
2	(a)	(i)	The H <sup>+</sup> ion in an (nitric) acid has been replaced by a metal ion <b>OR</b> by a Ca <sup>2+</sup> ion ✓	1	<b>DO NOT ALLOW</b> it has been produced by the reaction of an acid and a base as this is stated in the question.  <b>IGNORE</b> references to replacement by NH <sub>4</sub> <sup>+</sup> ions or positive ions. <b>ALLOW</b> H <b>OR</b> Hydrogen for H <sup>+</sup> ; <b>DO NOT ALLOW</b> Hydrogen atoms <b>ALLOW</b> Ca <b>OR</b> Calcium for Ca <sup>2+</sup> . <b>DO NOT ALLOW</b> Calcium atoms <b>ALLOW</b> 'metal' for 'metal ion'
		(ii)	2HNO <sub>3</sub> (aq) + Ca(OH) <sub>2</sub> (aq) → Ca(NO <sub>3</sub> ) <sub>2</sub> (aq) + 2H <sub>2</sub> O(l) Formulae ✓ Balance <b>AND</b> states ✓	2	<b>ALLOW</b> multiples <b>ALLOW</b> (aq) <b>OR</b> (s) for Ca(OH) <sub>2</sub>
		(iii)	Accepts a <b>proton</b> <b>OR</b> accepts H <sup>+</sup> ✓	1	<b>ALLOW</b> H <sup>+</sup> + OH <sup>-</sup> → H <sub>2</sub> O <b>ALLOW</b> OH <sup>-</sup> reacts with H <sup>+</sup> <b>OR</b> OH <sup>-</sup> takes H <sup>+</sup> <b>ALLOW</b> OH <sup>-</sup> 'attracts' H <sup>+</sup> if 'to form water' is seen  <b>DO NOT ALLOW</b> OH <sup>-</sup> neutralises H <sup>+</sup> ('neutralises' is in the question)
	(b)	(i)	Calculates correctly $\frac{0.0880 \times 25.0}{1000} = 2.20 \times 10^{-3}$ mol <b>OR</b> 0.00220 mol ✓	1	<b>ALLOW</b> 0.0022 <b>OR</b> $2.2 \times 10^{-3}$ mol
		(ii)	Calculates correctly $\frac{0.00220}{2} = 1.10 \times 10^{-3}$ mol <b>OR</b> 0.00110 mol ✓	1	<b>ALLOW</b> 0.0011 <b>OR</b> $1.1 \times 10^{-3}$ mol  <b>ALLOW</b> ECF for answer (i)/2 as calculator value or correct rounding to 2 significant figures or more but ignore trailing zeroes
		(iii)	$\frac{0.00110 \times 1000}{17.60} = 0.0625$ mol dm <sup>-3</sup> <b>OR</b> $6.25 \times 10^{-2}$ mol dm <sup>-3</sup> ✓	1	<b>ALLOW</b> 0.063 <b>OR</b> $6.3 \times 10^{-2}$ mol dm <sup>-3</sup>  <b>ALLOW</b> ECF for answer (ii) × 1000/17.60 <b>OR</b> ECF from (i) for answer (i)/2 × 1000/17.60 as calculator value or correct rounding to 2 significant figures or more but ignore trailing zeroes

	(c)	(i)	(The number of) Water(s) of crystallisation ✓	1	<b>IGNORE</b> hydrated <b>OR</b> hydrous
		(ii)	142.1 ✓  $x = \frac{(322.1 - 142.1)}{18.0} = 10$ ✓	2	<b>ALLOW</b> 142 <b>ALLOW</b> $M_r$ expressed as a sum  <b>ALLOW</b> ECF from incorrect $M_r$ and $x$ is <b>calculated correctly</b>  <b>ALLOW</b> ECF values of $x$ from nearest whole number to calculator value  <b>ALLOW</b> 2 marks if final answer is 10 <b>without any working</b>
			<b>Total</b>	<b>10</b>	


Question			Expected Answers	Marks	Additional Guidance
3	(a)	(i)	(Electrostatic) <b>attraction</b> between oppositely charged ions. ✓	1	<b>IGNORE</b> force <b>IGNORE</b> references to transfer of electrons <b>MUST</b> be ions, not particles
		(ii)	Mg shown with either 8 or 0 electrons <b>AND</b> S shown with 8 electrons <b>with</b> 2 crosses and 6 dots (or vice versa) ✓  Correct charges on both ions ✓  	2	Mark charges on ions and electrons independently <b>For first mark</b> , if 8 electrons are shown around the Mg then 'extra electrons' around S must match the symbol chosen for electrons around Mg  Shell circles not required  <b>IGNORE</b> inner shell electrons  Brackets are not required
	(b)	(i)	Electron pairs in covalent bonds shown correctly using dots and crosses in a molecule of the F <sub>2</sub> O ✓  Lone pairs correct on O and both F atoms ✓  	2	Must be 'dot-and-cross' circles for outer shells <b>NOT</b> needed <b>IGNORE</b> inner shells  Non-bonding electrons of O do not need to be shown as pairs  Non-bonding electrons of F do not need to be shown as pairs
		(ii)	Predicted bond angle 104–105°. ✓  There are 2 bonded pairs and 2 lone pairs ✓ Lone pairs repel more than bonded pairs ✓	3	<b>ALLOW</b> 103–105° (103° is the actual bond angle)  <b>ALLOW</b> responses equivalent to second marking point. e.g. There are 4 pairs of electrons and 2 of these are lone pairs <b>ALLOW</b> 'bonds' for 'bonded pairs' <b>DO NOT ALLOW</b> 'atoms repel' <b>DO NOT ALLOW</b> electrons repel <b>ALLOW</b> LP for 'lone pair' <b>ALLOW</b> BP for bonded pair <b>ALLOW</b> LP repel more if bonded pairs have already been mentioned

Question	Expected Answers	Marks	Additional Guidance
(c) (i)	<p>(At least) two <math>\text{NH}_3</math> molecules with correct dipole shown with at least one H with <math>\delta^+</math> and one N with <math>\delta^-</math> ✓</p> <p>(Only) one hydrogen bond from N atom on one molecule to a H atom on another molecule ✓</p> <p>Lone pair shown on the N atom and hydrogen bond must hit the lone pair ✓</p> 	3	<p><b>DO NOT ALLOW</b> first mark for ammonia molecules with incorrect lone pairs</p> <p><b>DO NOT ALLOW</b> first mark if <math>\text{H}_2\text{O}</math>, <math>\text{NH}_2</math> or <math>\text{NH}</math> is shown</p> <p><b>ALLOW</b> hydrogen bond need not be labelled as long as it clear the bond type is different from the covalent N–H bond</p> <p><b>ALLOW</b> a line (i.e. looks like a covalent bond) as long as it is labelled 'hydrogen bond'</p> <p><b>ALLOW</b> 2-D diagrams</p> <p><b>ALLOW</b> two marks if water molecules are used. One awarded for a correct hydrogen bond and one for the involvement of lone pair</p>
(ii)	<p>Liquid <math>\text{H}_2\text{O}</math> is denser than solid ✓</p> <p>In solid state <math>\text{H}_2\text{O}</math> molecules are held apart by hydrogen bonds <b>OR</b> ice has an open lattice ✓</p> <p><b>OR</b></p> <p><math>\text{H}_2\text{O}</math> has a relatively high boiling point <b>OR</b> melting point ✓</p> <p>(relatively strong) hydrogen bonds need to be broken <b>OR</b> a lot of energy is needed to overcome hydrogen bonds <b>OR</b> hydrogen bonds are strong ✓</p>	2	<p>ORA</p> <p><b>ALLOW</b> ice floats for first mark</p> <p><b>ALLOW</b> higher melting <b>OR</b> boiling point than expected</p> <p><b>DO NOT ALLOW</b> <math>\text{H}_2\text{O}</math> has a high melting / boiling point</p> <p><b>ALLOW</b> other properties caused by hydrogen bonding not mentioned within the specification</p> <p>E.g. high surface tension – strong hydrogen bonds on the surface</p>
	<b>Total</b>	<b>13</b>	

Question		Expected Answers	Marks	Additional Guidance
4	(a)	<p><i>Advantage</i> removes or kills bacteria <b>OR</b> kills germs <b>OR</b> kills micro-organisms <b>OR</b> make it safe to drink <b>OR</b> sterilises water <b>OR</b> disinfects water ✓</p> <p><i>Disadvantage</i> it is toxic <b>OR</b> poisonous <b>OR</b> could form chlorinated hydrocarbons ✓</p>	2	<p><b>ALLOW</b> to make water potable <b>IGNORE</b> virus <b>IGNORE</b> 'purifies water' <b>DO NOT ALLOW</b> 'antiseptic'</p> <p><b>ALLOW forms</b> carcinogens <b>OR</b> forms toxins <b>IGNORE</b> harmful <b>DO NOT ALLOW</b> 'it causes cancer' <b>DO NOT ALLOW</b> "It kills you"</p>
	(b)	$3d^{10} 4s^2 4p^5$ ✓	1	<p><b>ALLOW</b> <math>4s^2 3d^{10} 4p^5</math> <b>ALLOW</b> subscripts or <math>3D^{10}</math> <b>ALLOW</b> answers with <math>1s^2 2s^2 2p^6 3s^2 3p^6</math> appearing twice</p>
	(c) (i)	$Cl_2 + 2Br^- \rightarrow Br_2 + 2Cl^-$ ✓	1	<p><b>IGNORE</b> state symbols <b>ALLOW</b> any correct multiple including fractions</p>
	(ii)	Yellow / orange / red / brown ✓	1	<b>ALLOW</b> any combination of these, but no others
	(d) (i)	Disproportionation ✓	1	<p><b>ALLOW</b> versions which sound the same</p> <p><b>DO NOT ALLOW</b> disproportional <b>OR</b> disproportionate <b>OR</b> disproportion</p>
	(ii)	<p><math>Cl_2 + 2NaOH \rightarrow NaClO + NaCl + H_2O</math> ✓</p> <p><math>3Cl_2 + 6NaOH \rightarrow NaClO_3 + 5NaCl + 3H_2O</math></p> <p><math>Cl_2</math> and NaOH as reactants <b>AND</b> <math>NaClO_3</math> and NaCl as products ✓</p> <p>Rest of the equation ✓</p>	3	<p><b>ALLOW</b> multiples for either equation</p> <p><b>ALLOW</b> <math>3Cl_2 + 6NaOH \rightarrow 2NaClO_3 + 4NaCl + 3H_2</math></p>
	(iii)	$NaClO_4$ ✓	1	<b>ALLOW</b> $Na_3ClO_5$ etc
<b>Total</b>			<b>10</b>	



Question			Expected Answers	Marks	Additional Guidance
5	(a)	(i)	Potassium <b>AND</b> argon ✓	1	<b>ALLOW</b> K and Ar
		(ii)	They are arranged in increasing atomic number <b>OR</b> Neither would show properties <b>OR</b> trends of rest of group <b>OR</b> Neither would show properties <b>OR</b> trends of rest of period <b>OR</b> They are arranged by electron configuration ✓	1	<b>ALLOW</b> any correct property difference e.g. This would place a reactive metal in the same group as noble gases  <b>ALLOW</b> they do not fit in with the rest of the group
	(b)	(i)	$2\text{Mg} + \text{O}_2 \rightarrow 2\text{MgO}$ ✓	1	<b>ALLOW</b> multiples. Correct species must be seen <b>IGNORE</b> state symbols
		(ii)	Fizzes <b>OR</b> bubbles <b>OR</b> gas produced <b>OR</b> effervescing ✓  Mg dissolves <b>OR</b> Mg disappears <b>OR</b> a solution is formed ✓	2	<b>DO NOT ALLOW</b> 'carbon dioxide gas produced' <b>DO NOT ALLOW</b> 'hydrogen produced' without 'gas'  <b>ALLOW</b> 'it for Mg' <b>IGNORE</b> Mg reacts <b>IGNORE</b> temperature change <b>IGNORE</b> steam produced
		(iii)	Quicker <b>OR</b> more vigorous <b>OR</b> gets hotter	1	<b>MUST</b> be a comparison of a reaction observation, not just 'more reactive'  <b>ALLOW</b> any comparison of greater rate including more bubbles etc. <b>DO NOT ALLOW</b> more gas produced

Question	Expected Answers	Marks	Additional Guidance
(c)	<p>Mg has a <b>giant</b> structure ✓</p> <p>Mg has <b>metallic</b> bonding OR description of metallic bonding as positive ions and <b>delocalised</b> electrons ✓</p> <p>(There is electrostatic attraction between) positive ions and electrons ✓</p> <p>Cl has a simple molecular <b>OR</b> simple covalent (lattice) ✓</p> <p>Cl has van der Waals' forces (between molecules) <b>OR</b> Cl has instantaneous dipole–induced dipoles <b>OR</b> temporary dipole–temporary dipole ✓</p>	6	<p><b>Metallic OR delocalised</b> seen spelt correctly at least <b>ONCE</b></p> <p><b>DO NOT ALLOW</b> as label nuclei <b>OR</b> protons for positive ions</p> <p><b>ALLOW</b> labelled diagram of metallic bonding for second and third marks</p>  <p>Lattice must have at least two rows of positive ions. If a Mg ion is shown it must correct charge</p> <p><b>ALLOW</b> for labels: + ions, positive ions, cations</p> <p><b>DO NOT ALLOW</b> as label nuclei <b>OR</b> protons for positive ions</p> <p><b>ALLOW</b> e<sup>-</sup> or e as label for electron</p> <p><b>DO NOT ALLOW</b> '-' without label for electron</p> <p><b>Covalent OR molecule OR molecular</b> seen spelt correctly at least <b>ONCE</b></p> <p><b>ALLOW</b> Cl is a (covalent) <b>molecule</b></p> <p><b>IGNORE</b> Cl has intermolecular bonding</p>

		<p>van der Waals' forces are weak <b>and</b> metallic bonds are strong  <b>OR</b>  van der Waals' forces are weaker than metallic bonds  <b>OR</b>  Less energy is needed to overcome van der Waals' than metallic bonds ✓</p>		<p><b>ALLOW</b> ECF from incorrect descriptions of giant structure with strong bonds; e.g. Mg has giant ionic structure  <b>ALLOW</b> ECF from any incorrect intermolecular forces e.g. permanent dipole –dipole from marking point 5    <b>ALLOW</b> vdW easier to break  ORA</p>
(d)	(i)	<p>O goes from –2 to 0 ✓    N goes from +5 to +4 ✓    N is reduced <b>AND</b> O is oxidised ✓</p>	3	<p>Oxidation numbers may be seen with equation    Third mark is dependent upon seeing a reduction in oxidation number of N and an increase in oxidation number of O    <b>ALLOW</b> ECF for third mark for N is oxidised <b>and</b> O is reduced if incorrect oxidation numbers support this    <b>IGNORE</b> references to strontium  <b>IGNORE</b> references to electron loss <b>OR</b> gain    <b>DO NOT ALLOW</b> 'One increases and one decreases'</p>

	<b>(d)</b>	<b>(ii)</b>	<p>Calculates correctly:  Mol of <math>\text{Sr}(\text{NO}_3)_2 = \frac{5.29}{211.6} = 0.0250 \checkmark</math></p> <p>Calculates correctly:  Mol of gas = <math>5/2 \times 0.0250 = 0.0625 \checkmark</math></p> <p>Calculates correctly:  Volume of gas = <math>24.0 \times 0.0625 = 1.50 \text{ dm}^3 \checkmark</math></p>	<b>3</b>	<p><b>ALLOW</b> 0.025</p> <p><b>ALLOW</b> ECF for first answer <math>\times 2.5</math> as calculator value or correct rounding to 2 significant figures or more but ignore trailing zeroes</p> <p><b>ALLOW</b> ECF for second answer <math>\times 24(.0)</math> as calculator value or correct rounding to 2 significant figures or more but ignore trailing zeroes</p> <p><b>DO NOT ALLOW</b> ECF of first answer <math>\times 24(.0)</math> (which gives <math>0.6(0) \text{ dm}^3</math>) as this has not measured the volume of any gas, simply <math>0.0250 \text{ mol}</math> of solid <math>\text{Sr}(\text{NO}_3)_2</math> converted into a gas  i.e. This answer would give <b>one</b> mark</p> <p><b>ALLOW</b> <math>1.5 \text{ dm}^3</math></p> <p><b>ALLOW</b> ECF producing correct volume of <math>\text{NO}_2</math> only  i.e. <math>1.2(0) \text{ dm}^3</math> would give <b>two</b> marks</p> <p><b>OR</b></p> <p><b>ALLOW</b> ECF producing correct volume of <math>\text{O}_2</math> only  i.e. <math>0.3(0) \text{ dm}^3</math> would give <b>two</b> marks</p>
<b>Total</b>			<b>18</b>		