


**Q1.** Ammonia is manufactured by the Haber process in which the following equilibrium is established.



(a) Give **two** features of a reaction at equilibrium.

Feature 1 .....

.....

.....

.....

Feature 2 .....

.....

.....

.....

(2)

(b) Explain why a catalyst has no effect on the position of an equilibrium.

.....

.....

.....


.....

(Extra space) .....

.....

(2)

(c) The diagram shows how the equilibrium yield of ammonia varies with changes in pressure and temperature.



(i) Use the diagram to state the effect of an **increase** in pressure at constant temperature on the yield of ammonia. Use Le Chatelier's principle to explain this effect.

Effect on yield .....

Explanation .....

.....  
 .....  
 .....  
 .....  
 .....

(3)

(ii) Use the diagram to state the effect of an **increase** in temperature at constant pressure on the yield of ammonia. Use Le Chatelier's principle to explain this effect.

Effect on yield .....

Explanation .....

.....  
 .....  
 .....  
 .....

(3)

(d) At equilibrium, with a pressure of 35 MPa and a temperature of 600 K, the yield of ammonia is 65%.

(i) State why industry uses a temperature higher than 600 K.

.....  
.....

(1)

(ii) State why industry uses a pressure lower than 35 MPa.  
Do **not** include references to safety.

.....  
.....

(1)

(Total 12 marks)

**Q2.** The following dynamic equilibrium was established at temperature T in a closed container.



The value of  $K_c$  for the reaction was  $68.0 \text{ mol}^{-1} \text{ dm}^3$  when the equilibrium mixture contained 3.82 mol of P and 5.24 mol of R.

(a) Give the meaning of the term *dynamic equilibrium*.

.....  
.....  
.....  
.....  
*(Extra space)* .....

(2)

(b) Write an expression for  $K_c$  for this reaction.

.....  
.....

(1)

- (c) The volume of the container was 10.0 dm<sup>3</sup>.

Calculate the concentration, in mol dm<sup>-3</sup>, of **Q** in the equilibrium mixture.

.....  
.....  
.....  
.....  
.....  
.....

*(Extra space)* .....

.....

(4)

- (d) State the effect, if any, on the equilibrium amount of **P** of increasing the temperature.  
All other factors are unchanged.

.....

(1)

- (e) State the effect, if any, on the equilibrium amount of **P** of using a container of larger volume. All other factors are unchanged.

.....

(1)

- (f) State the effect, if any, on the value of **K<sub>c</sub>** of increasing the temperature.  
All other factors are unchanged.

.....

(1)

- (g) State the effect, if any, on the value of **K<sub>c</sub>** of using a container of larger volume.  
All other factors are unchanged.

.....

(1)


(h) Deduce the value of the equilibrium constant, at temperature  $T$ , for the reaction



.....  
 .....

(1)  
 (Total 12 marks)

**Q3.** The diagram below shows a Maxwell–Boltzmann distribution for a sample of gas at a fixed temperature.  $E_a$  is the activation energy for the decomposition of this gas.



(a) (i) On this diagram, sketch the distribution for the same sample of gas at a higher temperature. (2)

(ii) With reference to the Maxwell–Boltzmann distribution, explain why an increase in temperature increases the rate of a chemical reaction.

.....  
 .....  
 .....  
 .....  
 .....

(2)

- (b) Dinitrogen oxide (N<sub>2</sub>O) is used as a rocket fuel. The data in the table below show how the activation energy for the decomposition of dinitrogen oxide differs with different catalysts.



	E <sub>a</sub> / kJ mol <sup>-1</sup>
Without a catalyst	245
With a gold catalyst	121
With an iron catalyst	116
With a platinum catalyst	136

- (i) Use the data in the table to deduce which is the most effective catalyst for this decomposition.

.....

(1)

- (ii) Explain how a catalyst increases the rate of a reaction.

.....  
 .....  
 .....  
 .....

(2)

(Total 7 marks)

**Q4.** Hess's Law is used to calculate the enthalpy change in reactions for which it is difficult to determine a value experimentally.

- (a) State the meaning of the term *enthalpy change*.

.....

(1)


- (b) State Hess's Law.

.....  
 .....  
 .....

(1)

(c) Consider the following table of data and the scheme of reactions.

Reaction	Enthalpy change / kJ mol <sup>-1</sup>
HCl(g) → H <sup>+</sup> (aq) + Cl <sup>-</sup> (aq)	-75
H(g) + Cl(g) → HCl(g)	-432
H(g) + Cl(g) → H <sup>•</sup> (g) + Cl <sup>•</sup> (g)	+963



Use the data in the table, the scheme of reactions and Hess's Law to calculate a value for  $\Delta H_{\text{r}}$

.....

.....

.....

.....

(3)  
(Total 5 marks)

**Q5.** Fritz Haber, a German chemist, first manufactured ammonia in 1909. Ammonia is very soluble in water.

(a) State the strongest type of intermolecular force between one molecule of ammonia and one molecule of water.

.....

(1)

(b) Draw a diagram to show how one molecule of ammonia is attracted to one molecule of water. Include all partial charges and all lone pairs of electrons in your diagram.

(c) Phosphine ( $\text{PH}_3$ ) has a structure similar to ammonia.

In terms of intermolecular forces, suggest the main reason why phosphine is almost insoluble in water.

.....  
 .....

(1)

(Total 5 marks)

**Q6.** Compound **A** is an oxide of sulphur. At 415 K, a gaseous sample of **A**, of mass 0.304 g, occupied a volume of 127  $\text{cm}^3$  at a pressure of 103 kPa.

State the ideal gas equation and use it to calculate the number of moles of **A** in the sample, and hence calculate the relative molecular mass of **A**.

(The gas constant  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ )

*Ideal gas equation* .....

*Calculation* .....

.....  
 .....  
 .....  
 .....

(Total 5 marks)

**Q7.** Which change requires the largest amount of energy?



(Total 1 mark)



**Q8.** A saturated aqueous solution of magnesium hydroxide contains  $1.17 \times 10^{-3}$  g of  $\text{Mg}(\text{OH})_2$  in  $100 \text{ cm}^3$  of solution. In this solution, the magnesium hydroxide is fully dissociated into ions.

What is the concentration of  $\text{Mg}^{2+}(\text{aq})$  ions in this solution?

- A  $2.82 \times 10^{-2} \text{ mol dm}^{-3}$
- B  $2.01 \times 10^{-3} \text{ mol dm}^{-3}$
- C  $2.82 \times 10^{-3} \text{ mol dm}^{-3}$
- D  $2.01 \times 10^{-4} \text{ mol dm}^{-3}$

(Total 1 mark)

**Q9.** The table below shows some information about three hydrochloric acid solutions used to clean bricks and concrete.

Cleaner	Acid content by mass / %	Price per $25 \text{ dm}^3$ / £
Plattern Concrete Acid	24.0	14.39
Dub-Lit Brick Cleaner	28.9	16.99
Conpat Brick Acid	35.9	24.99

Use the data in the table above to determine the cleaner that offers the best value for money, based on acid content. Show your working.

.....

.....

.....

.....

(Total 1 mark)

**Q10.** Which of these pieces of apparatus has the lowest percentage uncertainty in the measurement shown?

- A** Volume of 25 cm<sup>3</sup> measured with a burette with an uncertainty of  $\pm 0.1$  cm<sup>3</sup>.
- B** Volume of 25 cm<sup>3</sup> measured with a measuring cylinder with an uncertainty of  $\pm 0.5$  cm<sup>3</sup>.
- C** Mass of 0.150 g measured with a balance with an uncertainty of  $\pm 0.001$  g.
- D** Temperature change of 23.2 °C measured with a thermometer with an uncertainty of  $\pm 0.1$  °C.

**(Total 1 mark)**